

NYMPHAEA Folia naturae Bihariae	XLVIII	109 - 130	Oradea, 2021
---	---------------	------------------	---------------------

Sporadic notes on the land snail (Gastropoda) fauna of the North Bihor Mountains, Romania

Tamás DOMOKOS¹ & Barna PÁLL-GERGELY²

¹H–1124 Budapest, Bűrök utca 24–26. E, II.4., e-mail: tamasdomokos@freemail.hu

²Agrártudományi Kutatóközpont, Növényvédelmi Intézet, ELKH, H–1022 Budapest, Herman Ottó út 15., e-mail: pallgergely2@gmail.com

Abstract: During a sum of fifteen-day fieldwork in 2007, 2008, 2009, 2010 and 2015 in the North Bihor Mountains, we collected 258 samples of 62 terrestrial and freshwater snail and a bivalve species. *Drobacia banatica*, *Ruthenica filograna*, *Strigillaria vetusta*, *Vitrina pelucida* were found to be relatively frequent. However, *Alopiä bielzi*, *Cochlodina marisi*, *Ena montana*, *Merdigera obscura*, *Euomphalia strigella*, *Fruticola fruticum*, *Laciniaria plicata*, *Orcula jetschini*, *Aspasita triaria* are considered rare.

Keywords: constancy, distribution curves, manual singling, mass sampling, number of individuals, Mollusc fauna

Introduction

The Bihar Mountains (Munții Bihorului) are the most prominent and highest part of the Apuseni Mountains in the Western Romanian Carpathians. That mountain range comprises of the Codru Moma Mountains in the west, the Pădurea Craiului Mountains in the northwest, the Vlădeasa in the northeast, and the Gilău Moun-

tains in the east. Its highest peak is Cucurbăta Mare (1849 m a.s.l.). The northern part mainly consists of karst-forming limestone, whereas the southern part consists of crystalline shales. The two subunits are separated by the Arieș and Crișul Băița Valleys.

The massif is part of Apuseni Natural Park, it is 25 km long from northwest to southeast, and has a width of 14 km. The area is characterized by an extraordinary diversity of karstic formations, such as caves, canyons and gorges. The Padiș–Cetățile Ponorului karstic plateau, with its 54 km² limestone outcrop, is a significant exokarst formation in Europe (Ponta 1998, Ponta & Onac 2018).

The annual rainfall in the Bihor Mountains reaches 1400 mm. The mean annual temperature is 2°C. The average in January is between -5 and -9 °C, and in July ranges between 8 and 16 °C. The Bihor Mountains are the source of the rivers Someșul Cald, Crișul Negru, Crișul Pietros and Arieș.

Although the species diversity is high, the area has not been investigated intensively yet. Data on malacofauna were published in monographs of Csiki (1918), Soós (1943) And Grossu (1981, 1983, 1987), and in some additional papers by various other authors (Mocsáry 1972, Lupu 1966, Varga 1983, Boeters, et al. 1989, Bába & Sárkány-Kiss 1999, 2001, Sîrbu & Benedek 2004, Domokos & Váncsa 2005, Sîrbu 2006, Váncsa 2006, Falniowski et al. 2009, Ferhér et al. 2009, Lengyel & Páll-Gergely 2010, Deli & Domokos 2011, Deli & Subai 2011, Domokos 2013, 2015, 2016, Farkas & Deli 2015, Domokos & Lennert 2016, Domokos et al. 2018).

Material and method

Snail specimens were collected by manual singling (S) and mass sampling (M – 3dm³). At one occasion, flotsam material of ca. 2 litres was collected from the Sighiștel brook. Apart from this, aquatic molluscs were only occasionally collected.

The collected shells were cleaned and dried, and were identified based on their morphology using the following sources: Soós (1943), Grossu (1981, 1983, 1987, 1993), Kerney et al. (1983), Richnovszky & Pintér (1979), Deli & Subai (2011), Nordsieck (2007), Welter-Schultes (2012).

The nomenclature of check list was composed following MolluscaBase (<http://www.molluscabase.org>). Shells are deposited in the private collection of the senior author.

Description of the sampling sites in chronological order

Between 2007 and 2015, 25 sites were sampled in Apuseni Mountains Natural Park:




Figure 1. Hydrographic sketch map of sampling sites in the North Bihor Mountains (see text).

1. Alba county, Gârda de Sus, Peștera Ghețarul de la Scărișoara, 1060 m a.s.l., western side of pothole entrance, detric rock covered by moss and forest litter (M), 5. IV. 2007.
2. Bihor county, Pietroasa, Vale Bulz, Left side of the brook, 540 m a.s.l., under bushes (S), 10. VI. 2007.
3. Bihor county, Pietroasa, Cetățile Ponorului, cave near Dolina 3., 1000 m a.s.l., moss and forest litter-covered cliffs facing north and south (M), 11. VI. 2007.
4. Bihor county, Cheile Someșului Cald, „Babilon kapuja”, 1250 m a.s.l., stones facing south, and ground covered by great burdock (S), 12. VI. 2007.
5. Bihor county, Pietroasa, gorge east of Cabana Cetățile Ponorului, 1050 m a.s.l., stones (S), 12. VI. 2007.
6. Bihor county, Pietroasa, Valea Seacă, rock wall west of Cabana Cetățile Ponorului, 1050 m a.s.l., forest litter and moss (S), 13. VI. 2007.
7. Bihor county, Pietroasa, Izbuc Ponor, 1140 m a.s.l., hill-side facing northeast, covered with decaying plant material (S), 13. VI. 2007.
8. Bihor county, Pietroasa, Valea Brădeșanului, 0.5 km N of the Izbuc Ponor, 1200 m a.s.l., the left side of the brook, great burdock-covered bank, (S), 13. V. 2007.
9. Bihor county, Sighiștel, Cheile Sighiștelui, around a cave-spring, 380 m a.s.l., rock ledges covered by forest litter and moss (M), 29. IX. 2007.
10. Bihor county, Pasul Vârtop, 1230 m a.s.l., 7 km W of the settlements Arieșeni, dell leading to Groapa Ruginoasa, great burdock near the current (S), 29. IX. 2007.
11. Bihor county, Chișcău, cliff wall near Peștera Urșilor (cave) (S) VII. 2008.
12. Bihor county, Pietroasa, Stațiunea Boga, downhill near Cascada Oșel (S), (leg. Sarkadi, L.) VIII. 2008.
13. Alba county, Gârda de Sus, Valea Ordâncușă, Poarta lui Ionel, ca. 800 m a.s.l., great burdock, moss (S) (leg. Sarkadi L.) X. 2008.
14. Bihor county, Chișcău, near ramp of the Peștera Urșilor (cave), 482 m a.s.l., hillside faces north, covered by rock-vegetation, occasionally with bushes, (M), 9. IV. 2009.
15. Bihor county, Chișcău, rock-glacier leading to the Peștera Urșilor (cave), in the vicinity of the ramp, 470 m a.s.l., (S). 9. IV. 2009.
16. Bihor county, Pietroasa, Valea Bulzului, Piatra Bulzului, 580 m a.s.l., south-exposed rock wall, woody and shady biotope with humid leaves (M), 1. XII. 2009.
17. Bihor county, Pietroasa, Stațiunea Boga, downhill near Peștera Dracului, humid leaf litter, (M), 1. XII. 2009.
18. Alba county, Gârda de Sus, Peștera Ghețarul de la Scărișoara, 1060 m a.s.l., western side of the path leading to the cave, detrital rock covered by moss and forest litter, (M), 25. IV. 2010.




Figure 2. Hydrographic sketch map of sampling sites in the North Bihor Mountains (see text).




Figure 3. Hydrographic sketch map of sampling sites in the North Bihor Mountains (see text).

19. Bihor county, Chișcău, Giulești, under Dealul Vârsecilor, Valea Seacă after confluence with Țiganul (brook), 800 m a.s.l., base of rock wall, beech forest, humid leaf litter (S), 6. XI, 2010.




Figure 4. Hydrographic sketch map of sampling sites in the North Bihor Mountains (see text).

20. Bihor county, Chișcău, Giulești, Valea Nucșoara, rock-glacier, 1000 m a.s.l., wet and shady biotope (M), 6. XI. 2010.

21. Bihor county, Sighiștel, Cheile Sighiștelului, banks of the brook at the beginning of the gorge (S), 7. XI. 2010.

22. Bihor county, Sighiștel, Cheile Sighiștelului, flotsam material collected at begin-




Figure 5. Hydrographic sketch map of sampling sites in the North Bihor Mountains (see text).

ning of gorge, 7. XI. 2010. 10. Quantity of the sample: 2 dm³.

23. Bihor county, Sighiștel, Cheile Sighiștelului, base of southeast-exposed rock wall at beginning of gorge, 380 m a.s.l., beech, hornbeam (M), 7. XI. 2010.

24. Bihor county, Sighiștel, Cheile Sighiștelului, base of northwest-exposed rock wall at beginning of gorge, 380 m a.s.l., beech, hornbeam (M), 7. XI. 2010.

25. Bihor county, Sighiștel, Cheile Sighiștelului, 600 m from end of gorge 600 m a.s.l., great burdock (M) 13. IX. 2015.

Results and discussion

Our investigations resulted in 62 species: 58 terrestrial snails, four freshwater snails and one bivalve. Altogether 258 samples were collected (see checklist and Tables 1./A, 1./B, 2/A, 2/B). Some rare and interesting species were found, such as *Alopias bielzi*, *Aspasita triaria*, *Cochlodina marisi*, *Ena montana*, *Merdigera obscura*, *Euomphalia strigella*, *Fruticicola fruticum*, *Laciniaria plicata*, and *Orcula jet-*




Figure 6. Hydrographic sketch map of sampling sites in the North Bihor Mountains (see text).

schini. *Drobacia banatica*, *Ruthenica filograna*, *Strigillaria vetusta* (Figs. 7–10.), and *Vitrina pellucida* are relatively frequent in the northern Bihor Mts.

Based on the collected specimens it could be stated that the following species are very variable in terms of shell morphology: *Drobacia banatica* (Fig.11.), *Granaria frumentum*, *Ruthenica filograna*, *Strigillaria vetusta* (Lengyel & Páll-Gergely 2010, Domokos 2013, 2016).

Annotated checklist of the molluscs of the Bihor Mountains

CLASS GASTROPODA

SUBCLASS PULMONATA

Aciculidae

Platyla banatica (Rossmässler, 1842)

Platyla microspira (Pini, 1884)

Platyla polita (W. Hartmann, 1840)

All these species are reported by Lengyel & Páll-Gergely (2010).

Of the three species only *P. microspira* was relatively frequent (constancy: 24%, number of collected individuals: 34), collected from Cheile Sigiștelului (380 m a.s.l.) and Peștera Ghețarul de la Scărișoara (1060 m a.s.l.).

Arionidae

Arion circumscriptus G. Johnston, 1828

In absence of living individual anatomical examination is not possible, therefore this species was identified on the basis of morphology and colour.

Camaenidae

Fruticicola fruticum (O. F. Müller, 1774)

Very rarely examined in the region by the present author. This species was collected only from rock-glacier in Valea Nucșoara.

Chondrinidae

Chondrina arcadica clienta (Westerlund, 1883)

Granaria frumentum (Draparnaud, 1801)

Clausiliidae

Alinda biplicata (Montagu, 1803)

It is a relatively rare species (number of collected individuals: five, constancy: 12,5%) here and elsewhere in Romania (Lengyel & Páll-Gergely 2010, Miklós Szekeres, pers. comm.).

Alinda stabilis (Pfeiffer, 1847)

Alopiä bielzii (Pfeiffer, 1849)

An isolated occurrence of the subspecies *Alopiä bielzi tenuis* (E.A. Bielz, 1861) was found on the south-southwestern cliffs of the Piatra Bulzului (Deli & Domokos 2011)

Cochlodina laminata (Montagu, 1803)

Cochlodina marisi (A. Schmidt, 1868)

Laciniaria plicata (Draparnaud, 1801)

Ruthenica filograna (Rossmässler, 1836)

This species is frequent in the Western Carpathian Mountains (Domokos 2015, Domokos & Lennert 2007, 2009, Domokos, Lennert & Márton 2010, Erőss




Figure 7. Number of individuals of clausiliid species in the Bihor Mountains.




Figure 8. Constancy of clausiliid species in the Bihor Mountains.

2015), and very variable in terms of shell morphology (Domokos & Lennert 2009, Domokos, Lennert & Márton 2010, Lengyel- Páll-Gergely 2010). *Ruthenica filograna* has the highest number of collected individuals (930) and constancy (68%).




Figure 9. Number of individuals of clausiliid species in the Bihor Mountains (based on Lengyel & Páll-Gergely 2010).




Figure10. Constancy of clausiliid species in the Bihor Mountains (based on Lengyel & Páll-Gergely 2010).

Strigillaria vetusta (Rossmässler, 1836).

A very variable species in terms of shell morphology (Lengyel & Páll-Gergely 2010). Similar to *Ruthenica filigrana*, it was found frequently (number of collected individuals: 376, constancy: 76%, see Figs. 7 and 8).




Figure 11. Frequency distribution histograms of width of *Drobacia banatica* in different biotopes (Sampling sites: 4 and 25).

Cochlicopidae

Cochlicopa lubrica (O. F. Müller, 1774)

Cochlicopa lubricella (Porro, 1838)

Ellobiidae

Carychium minimum O.F. Müller, 1774

Carychium tridentatum (Risso, 1826)

Enidae

Ena montana (Draparnaud, 1801)

This species was collected in the Ghețarul Scărișoara over 1000 m a.s.l.

Mastus bielzi (M. Kimakowicz, 1890)

Merdigera obscura (O. F. Müller, 1774)

Gastrodontidae

Zonitoides nitidus (O. F. Müller, 1774)

Helicidae

Caucasotachea vindobonensis (C. Pfeiffer, 1828)

Drobacia banatica (Rossmässler, 1838)

The largest specimen (width: 34.7 mm) was found in a rock-glacier (Nr.16. Chișcău, 09. 04. 2009.). In this sampling site the number of individuals was only five (arithmetic mean 34.2 mm)!

This is a very variable species, illustrated by specimens, which were collected at different elevations (Fig.11). *Drobacia banatica* is generalist and it has a large ecological plasticity and climatic variability. The number of individuals and constancy is illustrative of that (163 and 68%).

Arianta arbustorum (Linnaeus, 1758)

This species is found only in the alpine region of the Bihor Mountains (Cheile Someșului Cald – 1250 m, Vf. Boga –1340 m, Valea Brădetanului – 1200m).

Helix pomatia Linnaeus, 1758

Isognomostoma isognomostomos (Schröter, 1784)

A relatively rare species, its constancy reaches only 40%.

Faustina faustina (Rossmässler, 1835)

Hygromiidae

Euomphalia strigella (Draparnaud, 1801)

Kovacsia kovacsi (Varga & L. Pintér, 1972)

No living individuals were found, therefore the identification was based solely on shell morphology and on the ecology (rock-grassy, rock-shrub and open-wood).

Trochulus hispidus (Linnaeus, 1758)

Limacidae

Limax cinereoniger Wolf, 1803

Table 1/A: Sampling sites and number of species in Bihor Mts.

Species	Sampling sites												
	1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Acanthinula aculeata</i>													
<i>Aegopinella epipedostoma</i>				1		3						3	
<i>Aegopinella pura</i>	12		2					4	1				
<i>Alinda biplicata</i>													
<i>Alinda stabilis</i>						1	3	4				1	3
<i>Alopiia bielzi</i>													
<i>Arianta arbustorum</i>				4				5					
<i>Arion circumscriptus</i>			1										
<i>Aspasita triaria</i>	7								4				
<i>Carpathica calophana</i>			2	1				1		1			
<i>Carychium minimum</i>													
<i>Carychium tridentatum</i>	2												
<i>Caucasotachea vindobonensis</i>													
<i>Chondrina arcadica clienta</i>			1						17				
<i>Cochlicopa lubrica</i>													
<i>Cochlicopa lubricella</i>		1											
<i>Cochlodina laminata</i>	1					5	3		4				
<i>Cochlodina marisi</i>													
<i>Columella columella</i>													
<i>Drobacia banatica</i>	16	1		35	2		3	11	7	1		6	
<i>Ena montana</i>	3												
<i>Ena obscura</i>													
<i>Euomphalia strigella</i>									2		1		
<i>Faustina faustina</i>	14		1						9			3	
<i>Fruticicola fruticum</i>													
<i>Granaria frumentum</i>									37				
<i>Helix pomatia</i>			1				1						
<i>Isognomostoma isognomostomos</i>	10		3			1	5	1				2	
<i>Kovacsia kovacsi</i>		2							35				
<i>Laciniaria plicata</i>									2				
<i>Limax cinereoniger</i>			1										

Lymnaeidae

Galba truncatula (O. F. Müller, 1774)

Peregriana labiata (Rossmässler, 1835)

Table 1/B: Sampling sites and number of species in Bihor Mts.

Species	Sampling sites												
	1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Mastus bielzi</i>												2	
<i>Morlina glabra</i>													
<i>Orcula jetschini</i>			4										
<i>Oxychilus depressus</i>													
<i>Oxychilus draparnaudi</i>													
<i>Oxychilus montivagus</i>									6		1		
<i>Pyramidula pusilla</i>													
<i>Platyla banatica</i>	1		2										
<i>Platyla microspira</i>	4								7				
<i>Platyla polita</i>													
<i>Punctum pygmaeum</i>	5								1				
<i>Pupilla muscorum</i>													
<i>Ruthenica filograna</i>	10	18	48				11	7	138		4	6	6
<i>Sphyradium doliolum</i>													
<i>Strigillaria vetusta</i>	9	2	6	1	2	21	10		73		1	4	
<i>Trochulus bielzi</i>	5		3		1	8	5	1				10	
<i>Truncatellina cylindrica</i>									13				
<i>Vallonia costata</i>		2											
<i>Vallonia pulchella</i>									4				
<i>Vitrea contracta</i>									3				
<i>Vitrea botteri</i>													
<i>Vitrea crystallina</i>													1
<i>Vitrea diaphana</i>		1			1			13	4				1
<i>Vitrea jetschini</i>								1					
<i>Vitrea subrimata</i>			1										
<i>Vitrea szekeresi</i>	5	1	1					3					
<i>Vitrina pellucida</i>	6		14	1	2		6	5	10			1	
<i>Zonitoides nitidus</i>													
Number of species	16	8	16	6	5	6	9	12	20	2	4	10	4
Number of individuals	110	28	91	43	8	39	47	56	377	2	7	38	11
<i>Ancylus fluviatilis</i>													
<i>Bythiospeum</i> sp.													
<i>Galba truncatula</i>													
<i>Peregriana labiata</i>													
<i>Pisidium</i> sp.	–												

Table 2/A: Sampling sites and number of species in Bihor Mts. (C = constancy %, Σ = number of individuals)

Species	Sampling sites															Σ	C
	14	15	16	17	18	19	20	21	22	23	24	25					
<i>Acanthinula aculeata</i>	2								2	2	2					8	16
<i>Aegopinella epipedostoma</i>					5		1					1				14	24
<i>Aegopinella pura</i>	3			3	29				7		5					66	36
<i>Alinda biplicata</i>				3		1	1									5	12
<i>Alinda stabilis</i>					2				1							15	28
<i>Alopiia bielzi</i>			34													34	4
<i>Arianta arbustorum</i>																9	8
<i>Arion circumscriptus</i>																1	4
<i>Aspasita triaria</i>									5							16	12
<i>Carpathica calophana</i>																5	16
<i>Carychium minimum</i>	8						3									11	8
<i>Carychium tridentatum</i>	6			2		10			361							381	20
<i>Caucasotachea vindobonensis</i>	1									2						3	8
<i>Chondrina arcadica clienta</i>	34		24	3						85	39					203	28
<i>Cochlicopa lubrica</i>									12							12	4
<i>Cochlicopa lubricella</i>	3								2							6	12
<i>Cochlodina laminata</i>	9						2			4						28	28
<i>Cochlodina marisi</i>												1				1	4
<i>Columella columella</i>					9											9	4
<i>Drobacia banatica</i>		5	2	5	24	1	10		1			33				163	68
<i>Ena montana</i>																3	4
<i>Ena obscura</i>	3									1						4	8
<i>Euomphalia strigella</i>	1															4	12
<i>Faustina faustina</i>							2			4	6	3				42	32
<i>Fruticicola fruticum</i>							1									1	4
<i>Granaria frumentum</i>			2	1						91	49					180	20
<i>Helix pomatia</i>			1	1		1	1					1				7	28
<i>Isognomostoma isognomostomos</i>				5	1							1				29	36
<i>Kovacsia kovacsi</i>	32								5	51	30					155	24
<i>Laciniaria plicata</i>	3			2												7	12
<i>Limax cinereoniger</i>																1	4

Moitessieriidae

Bythiospeum sp.

„All these species were collected from fresh-water sources and mountain rivers in Western Carpathian Mountains and Banat.” (Grossu 1993. p. 297).

Table 2/B: Sampling sites and taxa in Bihor Mts. (C = constancy %, Σ = number of individuals)

Species	Sampling sites															Σ	C
	14	15	16	17	18	19	20	21	22	23	24	25					
<i>Mastus bielzi</i>						7							9	8			
<i>Morlina glabra</i>										3	3	1	7	12			
<i>Orcula jetschini</i>				4					1				9	12			
<i>Oxychilus depressus</i>									2				2	4			
<i>Oxychilus draparnaudi</i>				1									1	4			
<i>Oxychilus montivagus</i>	10		2	1			3						23	20			
<i>Pyramidula pusilla</i>	17		2				5			49	1		74	20			
<i>Platyla banatica</i>				2					5				10	16			
<i>Platyla microspira</i>					4				9	6	4		34	24			
<i>Platyla polita</i>					1								1	4			
<i>Punctum pygmaeum</i>	1				1	2			24				34	24			
<i>Pupilla muscorum</i>	1								16				17	8			
<i>Ruthenica filograna</i>	53		16		58	39	11			195	302	8	930	68			
<i>Sphyradium doliolum</i>	8					3	1			3	2		17	20			
<i>Strigillaria vetusta</i>	55		33	44	2	13	2			74	19	5	376	76			
<i>Trochulus bielzi</i>				4	3		1						41	40			
<i>Truncatellina cylindrica</i>	6								7	1	1		28	20			
<i>Vallonia costata</i>	2		1						11	4			20	20			
<i>Vallonia pulchella</i>	35								12				51	12			
<i>Vitrea contracta</i>	1												4	8			
<i>Vitrea botteri</i>				1									1	4			
<i>Vitrea crystallina</i>							1						2	8			
<i>Vitrea diaphana</i>	8				16						17		61	32			
<i>Vitrea jetschini</i>				4									5	8			
<i>Vitrea subrimata</i>	1												2	8			
<i>Vitrea szekeresi</i>				3									13	20			
<i>Vitrina pellucida</i>	7		2	5	53	2			1	5	13		133	64			
<i>Zonitoides nitidus</i>									1				1	4			
Number of species	26	1	11	19	14	10	15	–	19	17	15	9					
Number of individuals	310	5	119	94	208	79	45	–	485	580	492	54					
<i>Ancylus fluviatilis</i>								9	6				15				
<i>Bythiospeum</i> sp.									1				1				
<i>Galba truncatula</i>									5				5				
<i>Peregriana labiata</i>								14					14				
<i>Pisidium</i> sp.	1												1				

We collected one sample only from scum of the Sighiștel brook. According to Sirbu 2006, the following species can be found in the Pișolca and Coliboaia caves of Sighiștel Valley: *B. transsylvanicum* Rotarides, 1943, *B. carpathicum* Soós, 1940 and *P. leruthi* (C. Boettger, 1940).

Orculidae

Orcula jetschini (M. Kimakowicz, 1883)

Sphyradium doliolum (Bruguière, 1792)

Oxychilidae

Aegopinella epipedostoma (Fagot, 1879)

Welter-Schultes (2012) does not report this species from Romania, but Lengyel & Páll-Gergely (2010) listed it from the Bihor Mountains and published a drawing of its reproductive anatomy.

Aegopinella pura (Alder, 1830).

Carpathica calophana (Westerlund, 1881)

Morlina glabra (Rossmässler, 1838)

Oxychilus depressus (Sterki, 1880)

Oxychilus montivagus (M. Kimakowicz, 1890)

Planorbidae

Ancylus fluviatilis O.F. Müller, 1774

Punctidae

Punctum pygmaeum (Draparnaud, 1801)

Pupillidae

Pupilla muscorum (Linnaeus, 1758)

Pyramidulidae

Pyramidula rupestris (Draparnaud, 1801)

Spelaeodiscidae

Aspasita triaria (Rossmässler, 1839)

Collected only in Cheile Sighiștelui and Ghețarul Scărișoara.

Truncatellinidae

Truncatellina cylindrica (Férussac, 1807)

Valloniidae

Acanthinula aculeata (O. F. Müller, 1774)

Vallonia costata (O. F. Müller, 1774)

Vallonia pulchella (O. F. Müller, 1774)

Vertiginidae

Columella columella (G von Martens, 1830)

„Widespread in all the Carpathians, especially in the alpine zones, but also frequent in alluvia and löes” (Grossu 1993). This species became extinct in Pleistocene in Hungary (Fűköh et al. 1995).

Vitrinidae

Vitrina pellucida (O. F. Müller, 1774)

Zonitidae

Vitrea diaphana (S. Studer, 1820)

Vitrea contracta (Westerlund, 1871)

Vitrea crystallina (O. F. Müller, 1774)

Vitrea jetschini (M. Kimakowicz, 1890)

Vitrea subrimata (Reinhardt, 1871)

Vitrea szekeresi Deli & Subai 2011

The distribution of this species reaches its northern limit in the Bihor Mountains (Deli & Subai 2011).

CLASS BIVALVIA

SUBCLASS HETERODONTA

Sphaeriidae

Pisidium sp.

Only a single valve was collected from the flotsam of the Sighiștel brook.

Acknowledgements

This contribution is dedicated to the memory of Prof. Ioan Bușiția (1874–1953) from the high school of Beiuș, who dedicated all his active life to nature protection from the Apuseni Mountains. Prof. Bușiția noted on September 9 1931, in his touring guide-book of Apuseni Mountains [written by Czárán Gyula (1847-1906)]:

“and I wish from all my heart that the love for Nature brings many Hungarian tourists to us, so that we may delight in the beauty of our glades in brotherhood and good understanding, and practice tourism for the pleasure of our hearts and bodies”.

“și îmi doresc din toată inima, ca iubirea Naturii să aducă mulți turiști maghiari între noi, ca să putem să ne încântăm frumusețea poienilor noastre în fraternitate și în bună înțelegere, să practicăm turismul pentru plăcerea inimilor și a trupurilor noastre.”

“s óhajtom teljes szívemből, hogy a Természet iránti szeretet sok magyar turistát hozzon közibünk akikkel együtt gyönyörködhessünk havasaink szépségében, testvéri szeretettel, jó egyetértésben gyakoroljuk a turistaságot, szívünk gyönyörködtetésére s testeink edzésére.”

We are very grateful to Miklós Szekeres and Péter Subai for their help in the identification of Clausiliidae and Aciculidae, respectively. We thank to László Sarkadi, who put two samples (12.,13.) at my disposal.

References

- Bába, K., Sárkány-Kiss, A. 1999. - Contribution to the mollusc community of Someșul Cald/ Meleg Szamos gorges. *Tiscia, Monograph series* **3**: 275–278.
- Bába, K., Sárkány-Kiss, A. 2001. - The malacofauna of Bélavára – Scărița Belioara Mountain (Bihor Mountains, Romania). *Nachrichtenblatt der Ersten Vorarlberger Malakologischen Gesellschaft* **9**: 21–25.
- Boeters, H. D., Gittenberger, E., Subai, P. 1989. - Die Aciculidae (Mollusca: Gastropoda Prosobranchia). – *Zoologische Verhandlungen* **252**: 234.
- Csiki, E. 1918. - Mollusca, In: Fauna Regni Hungariae II. *Magyar Királyi Természettudományi Társulat* **6**: 1–44.
- Deli, T., Domokos, T. 2011. - An isolated occurrence of *Alopius bielzi tenuis* (E. A. Bielz, 1861) (Mollusca, Gastropoda, Clausiliidae) in the Bihor–Vlădeasa Mountains (Romania). *Nymphaea* **37**: 129–136.
- Domokos, T. 2013. - Shell morphological studies on *Granaria frumentum* (Draparnaud, 1801) populations from rockledges (Romania: Apuseni and Poiana Ruscă Mountains). *Nymphaea* **40**: 65–76.
- Domokos, T. 2015. - Sporadic data on the mollusc fauna of Rimetea and its environs (Trascău Mountains, Romania) with special reference to shell morphometry of *Alopius bielzi tenuis* (E. A. Bielz, 1861). *Nymphaea* **42**: 121–143.
- Domokos, T. 2016. - Is *Drobacia maeotica* a phantom species? Data on the shell morphology, genital morphology, and autecology of *Drobacia banatica* (Rossmässler, 1838) and *Drobacia maeotica* (Wenz, 1926), as well as some biophysical and biochemical notes. *Soosiana* **33**: 39–58.
- Domokos, T., Deli, T., Varga, A., Flaviu-Crișan, H., Balashov, I., Gheoca, V., Biatov, A., Szapapanos, B., Páll-Gergely, B. 2018. - Distribution of *Drobacia banatica* (Rossmässler, 1838) over time and space (Gastropoda: Stylommatophora: Helicidae). *Brukenenthal Acta Musei* **13**(3): 457–484.
- Domokos, T., Lennert, J. 2016. - Újabb adat a *Drobacia banatica* (Mollusca: Gastropoda: Helicidae) héjmorphológiájához. New data on the shell morphology of the *Drobacia banatica f. maeotica* (Mollusca)[Bazarul (Cetățile) Someșului Cald, Bihor Mts., Romania]. *Soosiana* **33**: 59–65.
- Domokos, T., Váncsa. 2005. - Data to mollusc fauna of Romania on the basis of the Munkácsy Mihály Múzeum's collection (Hungary, Békéscsaba). *Armonii naturale* **5**: 37–73.
- Falniowski, A., Szarovska, M., Sîrbu, I. 2009. - *Bythinella* Moquin-Tandon, 1856 (Gastropoda, Rissoidea: Bythinellidae) in Romania: its morphology with description of four new species. *Folia Malacologica* **17**(1): 21–36.
- Farkas, R., Deli, T. 2015. - *Chondrina tatrica* Ložek new species for the Romanian fauna (Gastropoda: Chondrinidae). *Malacologica Bohemoslovaca* **14**: 17–20.
- Grossu, A. V. 1981. - Gastropoda Romaniae, 3, Ed. Litera, București. 269 pp.
- Grossu, A. V. 1983. - Gastropoda Romaniae, 4, Ed. Litera, București. 563 pp.
- Grossu, A. V. 1987. - Gastropoda Romaniae, 2. Ed. Litera, București, 443 pp.
- Grossu, A. V. 1993. - The catalogue of the Molluscs from Romania. – *Travaux du Museum d'Istoire Naturelle „Grigore Antipa”* **33**: 291–366.

- Kerney, M. P., Cameron, R. A. D., Jungbluth, J. H. 1983. - Die Landschnecken Nord- und Mitteleuropas. P. Parey, Hamburg–Berlin. 384 pp.
- Lengyel, G. D., Páll-Gergely, B. 2010. - Notes on the landsnail (Gastropoda) and Harvestman (Opiliones) fauna of Bihor and Vlădeasa Mountains, Romania. – *Studii și Comunicări Seria Științele Naturii* **10-11**: 91–11.
- Lupu, D. 1966. - Un fossile vivant de la faune de la Romanie: *Chilostoma (Drobacia)* Wenz (Gastropoda–Pulmonata). *Travaux de Muséum d'Histoire Naturelle „Grigore Antipa”* **6**: 31–37.
- Mátyás, V. 1988. - Bihar-hegység turistakalauz. Sport. Budapest.
- Mocsáry, S. 1872. - Adatok Biharmegye faunájához. *Mathematikai és Természettudományi Közlemények* **10**: 163–200.
- Nordsiek, H. 2007. - Balea Gray 1824 and Alinda H & A. Adams are separated as genera (Gastropoda, Stylommatophora: Clausiliidae). *Mitteilungen der Deutschen Malakologischen Gesellschaft* **78**: 27–30.
- Ponta, G. 1998. - Karst of Romania 193–207. In: Global Karst Correlation (Eds: Daoxian, Y. & Zaihua, L.) Science Press and VSP BV. Beijing, New York.
- Ponta, G. M. L., Onac, B.P. (Eds.). 2018. Cave and Karst Systems of Romania. 544p, Springer.
- Richnovszky, A., Pintér, L. 1979. - A vízicsigák és kagylók (Mollusca) kishatározója. *Vízügyi Hidrobiológia* **6**: 1–206.
- Sîrbu, I. 2006. - The freshwater Mollusca from Crișana (Rivers basin, Romania) – *Travaux de Muséum d'Histoire Naturelle „Grigore Antipa”* **49**: 13–28.
- Sîrbu, I., Benedek, A. M. 2004. - The genus *Pisidium* C. Pfeiffer, 1821 (Bivalvia: Sphaeriidae) in the Romanian inner Carpathian Basin. *Travaux de Muséum d'Histoire Naturelle „Grigore Antipa”* **47**: 7–21.
- Soós, L. 1943. - A Kárpát-medence Mollusca-faunája. – In: Magyarország természetrajza, Állattani rész. Magyar Tudományos Akadémia, 478 pp.
- Váncsa, K. 2006. - A Magyar Természettudományi Múzeum Mollusca-gyűjteményében található Kovács Gyula-féle kollekció romániai vonatkozásai. *A Békés Megyei Múzeumok Közleményei* **28**: 41–60. [In Hungarian with English abstract]
- Varga, A. 1983. - Wirth Tibor Mollusca gyűjteménye. *Soosiana* **10/11**: 45–56.
- Welter-Schultes, F. 2012. - European non-marine molluscs, a guide for species identification. Planet Poster Editions.

Notă către autori

Nymphaea, Folia Naturae Bihariae, este anuarul Secției de Științele Naturii de la Muzeul Țării Crișurilor, Oradea, România. Contribuțiile științifice depuse pentru publicare vor fi redactate preferabil în limba engleză, sau unele articole din domeniul ocrotirii mediului, patrimoniului, conservare sau restaurare vor fi acceptate și în limba română, cu un rezumat (abstract) în limba engleză.

Vor fi considerate prioritare acele lucrări care abordează cercetarea naturii din zona Țării Crișurilor.

Înainte de acceptare, toate manuscrisele vor fi evaluate de comitetul de redacție și de referenți științifici. În acest scop, autorii sunt rugați să atașeze la manuscris o listă cu cel puțin 2 potențiali referenți în domeniu și adresa lor (e-mail).

Manuscrisele (pe un CD sau prin sistemul electronic) vor fi trimise pe adresa redacției:

Muzeul Țării Crișurilor

Str. Armatei Române 1/A, RO-410087 Oradea, Județul Bihor, România

E-mail: stiintelenaturii@mtariicrisurilor.ro

Copyright

Este o condiție esențială ca manuscrisele depuse să nu fi fost publicate în prealabil și nu vor apărea în alte publicații în mod simultan. Autorii sunt de acord ca, în cazul acceptării sau publicării lucrării, copyrightul acestuia să se transfere editurii.

Instrucțiuni de redactare

Textul manuscrisului va fi aranjat în felul următor: titlu; rezumat; cuvinte cheie (opțional); textul articolului integrând la final și eventualele mulțumiri (*Acknowledgements*) către persoane sau instituții; lucrări citate; tabele; figuri; explicația figurilor.

Titlul lucrării va fi scurt și informativ, cu caractere de 12 pct. boldate, centrat; prenumele și numele autorilor, cu caractere italice, vor fi însoțite de adresa lor (inclusiv e-mail).

Rezumatul (*Abstract*) informativ în limba engleză (plus în limba în care s-a redactat articolul) va fi sub forma unui singur paragraf, fără să depășească 3 % din lungimea totală. Acesta va cuprinde cele mai importante idei ale articolului, precum și concluziile (vor fi evitate prescurtările și citarea unor lucrări).

Textul integral al manuscrisului se va redacta la două rânduri (cu caractere Arial 10 pct.), aliniate la stânga, în formatul A4, lăsând pentru marginea dreaptă și stângă cel puțin 3 cm; toate paginile vor fi numerotate.

Lucrările citate vor fi trecute în ordine alfabetică (vor fi cuprinse numai lucrările citate în text). Se va folosi modelul următor:

Frost, DR. & Hillis, D. M. 1990. - Species in concept and practice: herpetological applications. *Herpetologica* **46**: 87-104.

Lindemann, R. & Yochelson, E. L. 1994. - Redescription of *Stylolina* (Incertae sedis). In: Landing, E. (Ed.): *Studies in Stratigraphy and paleontology*. New York State Museum, Bulletin 481: 149-160, New York.

Taylor, T.N. 1981. - *Palaeobotany: An introduction to fossil plant biology*. (McGraw-Hill Book Co.) New York, 589 p.

Danciu, M. 1974. - Studii geobotanice în sudul Muntelui Baraolt, Teză de doctorat, Universitatea din București, București, 175 p.

Tabelele vor fi anexate separat (*portrait* sau *landscape*, salvate în fișiere separate: ex. Table 1, 2, etc., preferabil Excel sau PDF), fiecare pe câte o foaie separată, nepaginată și vor fi prevăzute cu câte un cap de tabel. Acesta va conține numărul și titlul tabelului (dimensiunea maximă: 125x175 mm sau 175x125 mm).

Figurile se vor executa în tuș negru pe calc sau vor fi tipărite pe hârtie albă, de calitate foarte bună, fiecare pe foaie separată în format B 5 și numerotate consecutiv (Fig. 1, Fig. 2, etc.). Caracterele folosite după micșorarea figurii nu trebuie să fie mai mici de 2 mm. Vor fi acceptate figuri (diagrame, desene, fotografii) în formatul electronic (preferabil TIFF, EPS) cu dimensiunea maximă de 125x175 mm, cu o rezoluție minimă de 600 dpi pentru desene (*line drawings*) și 300 dpi pentru fotografii. Figurile generate de computer vor fi salvate în fișiere separate (Figure 1, 2, etc.). Locul preferat al figurilor se va marca în manuscris pe partea dreaptă a paginii respective.

Pentru fiecare articol publicat, autorii beneficiază de 15 extrase și de un volum al anuarului *Nymphaea*.